



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

A novel biomarker panel for triage of HPV positive women with high specificity for detection of clinically relevant cervical disease.

Citation for published version:

Bhatia, R, Stewart, J, Moncur, S, Cuschieri, K, Busby-Earle, C, Williams, A, Kavanagh, K & Howie, S 2017, 'A novel biomarker panel for triage of HPV positive women with high specificity for detection of clinically relevant cervical disease.', International Papillomavirus Conference 2017, Cape Town, South Africa, 28/02/17 - 4/03/17.

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



A novel biomarker panel for triage of HPV positive women with high specificity for detection of clinically relevant cervical disease

Ramya Bhatia¹, June Stewart², Sharon Moncur², Kate Cuschieri³, Camille Busby-Earle⁴, Alistair Williams⁴, Kim Kavanagh⁵, and Sarah Howie².

1. HPV Research Group, University of Edinburgh, UK, 2. Centre for Inflammation research, University of Edinburgh, UK, 3. Scottish HPV Reference Laboratory, Royal Infirmary of Edinburgh, UK, 4. Simpson Centre for Reproductive Health, Royal Infirmary of Edinburgh, UK, 5. Department of Mathematics and Statistics, Strathclyde University, UK.

Introduction

HR-HPV based primary cervical screening is now being implemented in several countries. Robust objective triage strategies for management of HR-HPV positive women are needed.

The objective of this study was to identify proteins involved in innate immunity at epithelial surfaces which could be used as biomarkers to risk stratify HR-HPV positive women.

Methods

Samples- Liquid based cytology (LBC) samples were obtained from both screening and colposcopy populations from the Scottish HPV archive.

HPV typing was done using the Abbott rtHPV test.

Protein extraction was done using chloroform isopropanol method (Canham *et al*, 2014).

Chemokine levels were measured using ProcartaPlex kit (Affymetrix eBioscience, UK) according to manufacturer's guidelines.

Results




Figure 1. Differences in chemokine proteins between CIN3 and all other groups. Pooled protein extracts from LBC samples (10mg of protein from each of 9 women) from women with different grades of disease and normal controls were analysed using chemokine proteomic array (R&D Systems, UK). **Left-** Each chemokine is printed in duplicate in the array. Patients with CIN3 showed a very different pattern of chemokine expression from any other group. **Right-** Expression level of chemokines in HR-HPV positive CIN3 relative to +ve control dots.




Figure 2. Six chemokines are upregulated in HR-HPV positive women with CIN2+. Results of a case control study of LBC samples from HR-HPV positive women with CIN2+ (cases, n=268) and cytology normal or CIN1 (controls, n=459) using ProcartaPlex kit. **Left-** Bar charts indicating levels of protein, *** = p<0.001 (Kruskal-Wallis test with Dunn's multiple comparison post-test). **Right-** Multivariate analysis of different predictive models to assess sensitivity (Sens), Specificity (Spec), Positive Predictive Value (PPV) and Negative Predictive Value (NPV) for detection of CIN2+.

Outcome

A panel of chemokines has been identified which could serve as an objective biomarker for HPV infection associated with CIN2+. This has potential as a triage strategy for HPV primary screening.